


Lecture I: Introduction of Methods and Concepts

January 2017



(c) Charley Harper, "Tree of Life"




Theoretical

Evolutionary

Ecology

(c) Charley Harper, "Tree of Life"




Theoretical


Evolutionary

Ecology


(c) Charley Harper, "Tree of Life"



(c) Charley Harper, "Tree of Life"



(c) Charley Harper, "Tree of Life"



(c) Charley Harper, "Tree of Life"

Seven steps to modelling a biological problem

[Otto and Day, 2007]

Seven steps to modelling a biological problem

1 Formulate the question

[Otto and Day, 2007]

Seven steps to modelling a biological problem

1 Formulate the question

What do you want to know?

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients

[more soon]

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system

Life-cycle, flow diagram

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system

Write down the equations

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system
- 5 Analyze the equations

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system
- 5 Analyze the equations
 - ▶ *Graphical analysis*
 - ▶ *Simulations*
 - ▶ *Equilibrium and stability analyses*
 - ▶ *Deriving general solutions*
 - ▶ *Determining long-term or asymptotic behavior*

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system
- 5 Analyze the equations
- 6 Checks and balances

[Otto and Day, 2007]


Seven steps to modelling a biological problem

- 1 Formulate the question
- 2 Determine basic ingredients
- 3 Qualitatively describe the system
- 4 Quantitatively describe the system
- 5 Analyze the equations
- 6 Checks and balances

Generality, alternative models

[Otto and Day, 2007]

Seven steps to modelling a biological problem

- 1 Formulate the question
 - 2 Determine basic ingredients
 - 3 Qualitatively describe the system
 - 4 Quantitatively describe the system
 - 5 Analyze the equations
 - 6 Checks and balances
 - 7 Relate results back to original question
- 

[Otto and Day, 2007]

Outline

Introduction

Defining a model

Basic ingredients

Features of evolutionary models

Discrete traits

Quantitative traits

Basic ingredients of a model

Variables

- ▶ Define the variables

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables

e.g., $p_i(t)$, frequency of individuals of type i at time t

$N_j(z, t)$, number of individuals of type z in deme j at time t

$X_j(t)$, indicator of whether there is a mutant at site j at time t

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables

e.g., $p_i(t)$, frequency of individuals of type i at time t

$N_j(z, t)$, number of individuals of type z in deme j at time t

$X_j(t)$, indicator of whether there is a mutant at site j at time t

- ▶ Identify constraints on the variables

e.g., $0 \leq p \leq 1$

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables
- ▶ Identify constraints on the variables

Time

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables
- ▶ Identify constraints on the variables

Time

- ▶ Is time continuous or discrete?

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables
- ▶ Identify constraints on the variables

Time

- ▶ Is time continuous or discrete?
- ▶ What is the time scale?

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables
- ▶ Identify constraints on the variables

Time

- ▶ Is time continuous or discrete?
- ▶ What is the time scale?

Parameters

[Otto and Day, 2007]

Basic ingredients of a model

Variables

- ▶ Define the variables
- ▶ Identify constraints on the variables

Time

- ▶ Is time continuous or discrete?
- ▶ What is the time scale?

Parameters


- ▶ Define the model's parameters;
- ▶ Identify constraints.

[Otto and Day, 2007]

Some features of evolutionary models

Reproduction

Clonal or sexual, with or without mate choice.




[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models

Reproduction

Clonal or sexual, with or without mate choice.



[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models



(c) Wikimedia



(c) Wikimedia




(c) Wikimedia

Reproduction

Clonal or sexual, with or without mate choice.

Evolutionary forces

- ▶ Drift
- ▶ Selection
- ▶ Migration
- ▶ Mutation



[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models



(c) Wikimedia



(c) Wikimedia




(c) Wikimedia

Reproduction

Clonal or sexual, with or without mate choice.

Evolutionary forces

- ▶ Drift
- ▶ Selection
- ▶ Migration
- ▶ Mutation



[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models



(c) Wikimedia



(c) Wikimedia




(c) Wikimedia

Reproduction

Clonal or sexual, with or without mate choice.

Evolutionary forces

- ▶ Drift
- ▶ Selection
- ▶ Migration
- ▶ Mutation



[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models



(c) Wikimedia



(c) Wikimedia




(c) Wikimedia

Reproduction

Clonal or sexual, with or without mate choice.

Evolutionary forces

- ▶ Drift
- ▶ Selection
- ▶ Migration
- ▶ Mutation



[Fisher, 1930, Wright, 1931]; F. Massol

Some features of evolutionary models



(c) Wikimedia



(c) Wikimedia




(c) Wikimedia

Reproduction

Clonal or sexual, with or without mate choice.


Evolutionary forces

- ▶ Drift
- ▶ Selection
- ▶ Migration
- ▶ Mutation



[Fisher, 1930, Wright, 1931]; F. Massol

Some other features...



... that we won't consider here.

[Laland et al., 2015]

A key concept: fitness

This survival of the fittest [...] is that which Mr. Darwin has called 'natural selection, or the preservation of favoured races in the struggle for life.'

Herbert Spencer, *The principles of biology* (1864)

A key concept: fitness

This survival of the fittest [...] is that which Mr. Darwin has called 'natural selection, or the preservation of favoured races in the struggle for life.'

Herbert Spencer, *The principles of biology* (1864)

What is “fitness”?

A key concept: fitness

This survival of the fittest [...] is that which Mr. Darwin has called 'natural selection, or the preservation of favoured races in the struggle for life'

Herbert Spencer, *The principles of biology* (1864)

What is “fitness”?

not this:

Dear Dr. Florence Débarre,
Warm Greetings from [Journal of Aerobics and Fitness](#).
We greatly appreciate your valuable contribution towards the scientific community.
We would like to invite you to contribute 1 page **Letter to Editor (Research, Review, Case Reports, Short Commentary of 1or 2**

Outline

Introduction

Defining a model

Discrete traits

Discrete time

Continuous time

Fitness components

Quantitative traits


Discrete traits – I) Discrete generations

Geometric population growth

$$A_t = (1 + a)^t A_0$$

$$B_t = (1 + b)^t B_0$$

$a = 0.41$ and $b = 0.26$.



[Hartl and Clark, 2007]

Discrete traits – I) Discrete generations

Geometric population growth



$$A_t = (1 + a)^t A_0$$

$$B_t = (1 + b)^t B_0$$

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = \left(\frac{1 + a}{1 + b} \right)^t \frac{A_0}{B_0}$$

$a = 0.41$ and $b = 0.26$.



[Hartl and Clark, 2007]

Discrete traits – I) Discrete generations

Geometric population growth

$$A_t = (1 + a)^t A_0$$



$$B_t = (1 + b)^t B_0$$

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

Relative fitness

$a = 0.41$ and $b = 0.26$.



[Hartl and Clark, 2007]

Discrete traits – I) Discrete generations

Geometric population growth

$$A_t = (1 + a)^t A_0$$



$$B_t = (1 + b)^t B_0$$

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

Relative fitness ★

$a = 0.41$ and $b = 0.26$.




[Hartl and Clark, 2007]

Discrete traits – I) Discrete generations (*continued*)

Proportion of A individuals

$$p_t = \frac{A_t}{A_t + B_t}$$




Discrete traits – I) Discrete generations (*continued*)

Proportion of A individuals

$$p_t = \frac{A_t}{A_t + B_t}$$

Change in the proportion of A individuals:

$$\Delta p_t = \frac{(w - 1)p_t(1 - p_t)}{wp_t + 1 - p_t}$$




Discrete traits – I) Discrete generations (*continued*)

Proportion of A individuals

$$p_t = \frac{A_t}{A_t + B_t}$$

Change in the proportion of A individuals:

$$\Delta p_t = \frac{(w - 1)p_t(1 - p_t)}{wp_t + 1 - p_t}$$



$$\begin{aligned}\Delta p_t &= p_{t+1} - p_t \\ &= \frac{(1 + a)A_t}{(1 + a)A_t + (1 + b)B_t} - p_t \\ &= \frac{wp_t}{wp_t + 1 - p_t} - p_t = \frac{(w - 1)p_t(1 - p_t)}{\bar{w}_t}\end{aligned}$$


Discrete traits – I) Discrete generations (*continued*)

Proportion of A individuals

$$p_t = \frac{A_t}{A_t + B_t}$$

Change in the proportion of A individuals:

$$\Delta p_t = \frac{(w - 1)p_t(1 - p_t)}{\bar{w}}$$



$$\begin{aligned}\Delta p_t &= p_{t+1} - p_t \\ &= \frac{(1+a)A_t}{(1+a)A_t + (1+b)B_t} - p_t \\ &= \frac{wp_t}{wp_t + 1 - p_t} - p_t = \frac{(w-1)p_t(1-p_t)}{\bar{w}_t}\end{aligned}$$

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Ratio of the number of A individuals to B individuals

$$\frac{A(t)}{B(t)} = \frac{A_0}{B_0} \exp((\alpha - \beta) t)$$

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Ratio of the number of A individuals to B individuals

$$\frac{A(t)}{B(t)} = \frac{A_0}{B_0} \exp(m t)$$

Relative fitness

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Ratio of the number of A individuals to B individuals

$$\frac{A(t)}{B(t)} = \frac{A_0}{B_0} \exp(m t)$$

Relative fitness ★

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Ratio of the number of A individuals to B individuals

$$\frac{A(t)}{B(t)} = \frac{A_0}{B_0} \exp(m t)$$

Relative fitness

Change in the proportion of A individuals

$$\frac{dp(t)}{dt} = (\alpha - \beta) p(t)(1 - p(t)).$$

Discrete traits – II) Continuous Time

Exponential population growth

$$A(t) = A_0 \exp(\alpha t)$$

$$B(t) = B_0 \exp(\beta t)$$

Ratio of the number of A individuals to B individuals

$$\frac{A(t)}{B(t)} = \frac{A_0}{B_0} \exp(m t)$$

Relative fitness

Change in the proportion of A individuals

$$\frac{dp(t)}{dt} = mp(t)(1 - p(t)).$$

Comparing discrete and continuous time

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$


$$\frac{A(t)}{B(t)} = \exp(m t) \frac{A_0}{B_0}$$

Comparing discrete and continuous time

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

Darwinian fitness



$$\frac{A(t)}{B(t)} = \exp(mt) \frac{A_0}{B_0}$$

Malthusian fitness




Comparing discrete and continuous time

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

Darwinian fitness



$$\frac{A(t)}{B(t)} = \exp(mt) \frac{A_0}{B_0}$$

Malthusian fitness



$$m = \ln(w) \approx w - 1$$

(when w is close to 1).

Comparing discrete and continuous time

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

$$\frac{A(t)}{B(t)} = \exp(mt) \frac{A_0}{B_0}$$

$$m = \ln(w) \approx w - 1$$

(when w is close to 1).

Change in the frequency of A individuals

$$\Delta p = \frac{(w - 1) p (1 - p)}{wp + 1 - p}$$

$$\frac{dp}{dt} = m p (1 - p).$$

Comparing discrete and continuous time

Ratio of the number of A individuals to B individuals

$$\frac{A_t}{B_t} = w^t \frac{A_0}{B_0}$$

$$\frac{A(t)}{B(t)} = \exp(mt) \frac{A_0}{B_0}$$

$$m = \ln(w) \approx w - 1$$

(when w is close to 1).

Change in the frequency of A individuals

$$\Delta p = \frac{(w-1)p(1-p)}{wp+1-p}$$

$$\approx (w-1)p(1-p)$$

(when w is close to 1).

$$\frac{dp}{dt} = mp(1-p).$$

Fitness components

Life stages at which selection can operate

[Bundgaard and Christiansen, 1972, Hedrick, 2011], with modifications

Fitness components

Life stages at which selection can operate

Fecundity selection

- ▶ Egg/seed production
- ▶ Sperm/pollen production

Viability selection

- ▶ Zygotic survival
- ▶ Rate of development
- ▶ Maternal-fetal interactions
- ▶ Sex-dependent survival

Sexual selection

- ▶ Assortative mating
- ▶ Male vigor
- ▶ Female choice

Gametic selection

- ▶ Nonrandom union of gametes
- ▶ Meiotic drive
- ▶ Self-incompatibility

[Bundgaard and Christiansen, 1972, Hedrick, 2011], with modifications

Fitness components

Life stages at which selection can operate

Fecundity selection

- ▶ Egg/seed production
- ▶ Sperm/pollen production
- ▶ Offspring production

Viability selection

- ▶ Zygotic survival
- ▶ Rate of development
- ▶ Maternal-fetal interactions
- ▶ Sex-dependent survival

Sexual selection

- ▶ Assortative mating
- ▶ Male vigor
- ▶ Female choice

Gametic selection

- ▶ Nonrandom union of gametes
- ▶ Meiotic drive
- ▶ Self-incompatibility

[Bundgaard and Christiansen, 1972, Hedrick, 2011], with modifications

Outline

Introduction

Defining a model

Discrete traits

Quantitative traits
Adaptive dynamics

Evolution of quantitative traits



vs.



Evolution of quantitative traits



Methods

- ▶ “Raw” model

No assumptions on the distribution.

e.g., use of integro-differential equations.

Evolution of quantitative traits



Methods

- ▶ “Raw” model

No assumptions on the distribution.

e.g., use of integro-differential equations.

- ▶ Quantitative genetics

Moment-based approaches

Evolution of quantitative traits



Methods

- ▶ “Raw” model
 - No assumptions on the distribution.*
 - e.g., use of integro-differential equations.*
- ▶ Quantitative genetics
 - Moment-based approaches*
- ▶ Adaptive dynamics

Evolution of quantitative traits



Methods

- ▶ “Raw” model
 - No assumptions on the distribution.*
 - e.g., use of integro-differential equations.*
- ▶ Quantitative genetics
 - Moment-based approaches*
- ▶ Adaptive dynamics

Adaptive dynamics (AD)

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

Adaptive dynamics (AD)

Assumptions

- ▶ Mutations are rare,

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

Adaptive dynamics (AD)

Assumptions

- ▶ Mutations are rare,

→ *Describe the evolution of a trait as a sequence of trait substitutions.*

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

Adaptive dynamics (AD)

Assumptions

- ▶ Mutations are rare,
→ *Describe the evolution of a trait as a sequence of trait substitutions.*
- ▶ Mutations are of small effect.

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

Adaptive dynamics (AD)

Assumptions

- ▶ Mutations are rare,
→ *Describe the evolution of a trait as a sequence of trait substitutions.*
- ▶ Mutations are of small effect.
→ *Local stability analysis*

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

Adaptive dynamics (AD)

Assumptions

- ▶ Mutations are rare,
→ *Describe the evolution of a trait as a sequence of trait substitutions.*
- ▶ Mutations are of small effect.
→ *Local stability analysis*

A Key Concept: Invasion fitness

Tells us whether an initially rare mutant can invade.

[Metz et al., 1992, Geritz et al., 1998, Doebeli, 2011]

AD: Example

Model

- ▶ Lotka-Volterra dynamics:

Carrying capacity

There is an optimal value of the trait, x_{opt}

$$K(x) = K_0 \exp\left(-\frac{(x - x_{opt})^2}{2v_K}\right),$$

Interaction term

Strongest when the traits are closest.

$$\alpha(x, y) = 1 - (x - y)^2.$$

- ▶ Mutation

Unbiased, of small variance.

$$\mu(x, z) = \frac{1}{\sigma_\mu \sqrt{2\pi}} \exp\left(-\frac{(x - z)^2}{2\sigma_\mu^2}\right).$$

Full model

$$\frac{\partial N(x, t)}{\partial t} = \left(1 - \frac{\int \alpha(x, y) N(y, t)}{K(x)} \right) N(x, t) - \int \mu(x, z) dz N(x, t) + \int \mu(z, x) N(z, t) dz.$$

AD: Example (3)

When mutation is very rare...

“Resident” (with trait x_r) vs. “mutant” (with trait x_m)

- ▶ Resident alone

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t)}{K(x_r)}\right) N_r(t).$$

AD: Example (3)

When mutation is very rare...

“Resident” (with trait x_r) vs. “mutant” (with trait x_m)

- ▶ Resident alone

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t)}{K(x_r)}\right) N_r(t).$$
$$\tilde{N}_r = K(x_r).$$

AD: Example (3)

When mutation is very rare...

“Resident” (with trait x_r) vs. “mutant” (with trait x_m)

- ▶ Resident alone

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t)}{K(x_r)}\right) N_r(t).$$
$$\tilde{N}_r = K(x_r).$$

- ▶ Resident + Mutant

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t) + \alpha(x_r, x_m)N_m(t)}{K(x_r)}\right) N_r(t),$$
$$\frac{dN_m(t)}{dt} = \left(1 - \frac{N_m(t) + \alpha(x_m, x_r)N_r(t)}{K(x_m)}\right) N_m(t).$$

AD: Example (3)

When mutation is very rare...

“Resident” (with trait x_r) vs. “mutant” (with trait x_m)

► Resident alone

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t)}{K(x_r)}\right) N_r(t).$$
$$\tilde{N}_r = K(x_r).$$

► Resident + rare Mutant

$$\frac{dN_r(t)}{dt} = 0,$$
$$\frac{dN_m(t)}{dt} = \left(1 - \frac{N_m(t) + \alpha(x_m, x_r)\tilde{N}_r}{K(x_m)}\right) N_m(t).$$

AD: Example (3)

When mutation is very rare...

“Resident” (with trait x_r) vs. “mutant” (with trait x_m)

► Resident alone

$$\frac{dN_r(t)}{dt} = \left(1 - \frac{N_r(t)}{K(x_r)}\right) N_r(t).$$
$$\tilde{N}_r = K(x_r).$$


► Resident + Mutant

$$\frac{dN_r(t)}{dt} = 0,$$
$$\frac{dN_m(t)}{dt} = \left(1 - \frac{N_m(t) + \alpha(x_m, x_r)\tilde{N}_r}{K(x_m)}\right) N_m(t).$$

Invasion fitness $w(x_m, x_r)$


Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.



Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\left. \frac{\partial w(x_m, x_r)}{\partial x_m} \right|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\left. \frac{\partial w(x_m, x_r)}{\partial x_m} \right|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\left. \frac{\partial w(x_m, x_r)}{\partial x_m} \right|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\left. \frac{\partial w(x_m, x_r)}{\partial x_m} \right|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\frac{\partial w(x_m, x_r)}{\partial x_m} \Big|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\left. \frac{\partial w(x_m, x_r)}{\partial x_m} \right|_{x_m=x_r}}_{\text{Selection gradient } D(x_r)}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\frac{\partial w(x_m, x_r)}{\partial x_m} \Big|_{x_m=x_r}}_{=D(x_r)=0}$$

Pairwise Invasibility Plots (PIP)

Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\frac{\partial w(x_m, x_r)}{\partial x_m} \Big|_{x_m=x_r}}_{=D(x_r)=0} + \frac{(x_m - x_r)^2}{2} \frac{\partial^2 w(x_m, x_r)}{\partial x_m^2} \Big|_{x_m=x_r}.$$

Pairwise Invasibility Plots (PIP)


Contour-plot of $w(x_m, x_r)$, showing regions where the mutant can invade.




$$w(x_m, x_r) \approx$$

$$\underbrace{w(x_r, x_r)}_{=0} + (x_m - x_r) \underbrace{\frac{\partial w(x_m, x_r)}{\partial x_m} \Big|_{x_m=x_r}}_{=D(x_r)=0} + \frac{(x_m - x_r)^2}{2} \frac{\partial^2 w(x_m, x_r)}{\partial x_m^2} \Big|_{x_m=x_r}.$$

Pairwise Invasibility Plots (continued)




Pairwise Invasibility Plots (continued)




(c) [Geritz et al., 1998]

Pairwise Invasibility Plots (continued)




(c) [Geritz et al., 1998]

Pairwise Invasibility Plots (continued)




(c) [Geritz et al., 1998]

Pairwise Invasibility Plots (continued)



(c) [Geritz et al., 1998]

Pairwise Invasibility Plots (continued)



Back to the example

Adaptive dynamics

- ▶ Singular strategy: $D(x^*) = 0$ when $x^* = x_{\text{opt}}$.

Back to the example

Adaptive dynamics

- ▶ Singular strategy: $D(x^*) = 0$ when $x^* = x_{\text{opt}}$.
- ▶ Convergence stability: $\frac{dD(x)}{dx} = -\frac{1}{v_K} < 0$.

Back to the example

Adaptive dynamics

► Singular strategy: $D(x^*) = 0$ when $x^* = x_{\text{opt}}$.

► Convergence stability: $\frac{dD(x)}{dx} = -\frac{1}{v_K} < 0$.

► Uninvadability (/ Evolutionary stability): $\left. \frac{\partial^2 w(x_m, x_r)}{\partial x_m^2} \right|_{x_m=x_r} = 2 - \frac{1}{v_K}$

Back to the example



Adaptive dynamics

- ▶ Singular strategy: $D(x^*) = 0$ when $x^* = x_{\text{opt}}$.

- ▶ Convergence stability: $\frac{dD(x)}{dx} = -\frac{1}{v_K} < 0$.

- ▶ Uninvadability (/ Evolutionary stability): $\left. \frac{\partial^2 w(x_m, x_r)}{\partial x_m^2} \right|_{x_m=x_r} = 2 - \frac{1}{v_K}$

Full model ($v_K = 0.2$)



Back to the example



Adaptive dynamics

- ▶ Singular strategy: $D(x^*) = 0$ when $x^* = x_{\text{opt}}$.

- ▶ Convergence stability: $\frac{dD(x)}{dx} = -\frac{1}{v_K} < 0$.

- ▶ Uninvadability (/ Evolutionary stability): $\left. \frac{\partial^2 w(x_m, x_r)}{\partial x_m^2} \right|_{x_m=x_r} = 2 - \frac{1}{v_K}$

Full model ($v_K = 0.55$)



Some take-home messages

Some dangers

- ▶ How do the model's results depend on the assumptions made?
- ▶ Do the results make sense biologically?

[Otto and Day, 2007]

Some take-home messages

Some dangers

- ▶ How do the model's results depend on the assumptions made?
- ▶ Do the results make sense biologically?

And ultimately...

- ▶ Can you test the model with real data / experiments?
- ▶ How does the model help us better understand and interpret the world?

[Otto and Day, 2007]

References

- Bundgaard, J. and Christiansen, F. B. (1972). Dynamics of polymorphisms. I. Selection components in an experimental population of *Drosophila melanogaster*. *Genetics*, 71(3):439.
- Doebeli, M. (2011). *Adaptive Diversification*. Princeton University Press.
- Fisher, R. A. (1930). *The genetical theory of natural selection*. Oxford University Press.
- Frank, S. (2011–2013). *Natural Selection Series*. Journal of Evolutionary Biology.
- Geritz, S., Kisdi, E., Meszéna, G., and Metz, J. (1998). Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evolutionary ecology*, 12(1):35–57.
- Hartl, D. L. and Clark, A. G. (2007). *Principles of population genetics*. Sinauer Associates, fourth edition.
- Hedrick, P. W. (2011). *Genetics of populations*. Jones & Bartlett Learning.
- Laland, K. N., Uller, T., Feldman, M. W., Sterelny, K., Müller, G. B., Moczek, A., Jablonka, E., and Odling-Smee, J. (2015). The extended evolutionary synthesis: its structure, assumptions and predictions. *Proceedings of the Royal Society of London B: Biological Sciences*, 282(1813).
- Levins, R. (1966). The strategy of model building in population biology. *American scientist*, 54(4):421–431.
- Metz, J. A., Nisbet, R. M., and Geritz, S. A. (1992). How should we define ‘fitness’ for general ecological scenarios? *Trends in Ecology & Evolution*, 7(6):198–202.
- Otto, S. P. and Day, T. (2007). *A Biologist’s Guide to Mathematical Modeling in Ecology and Evolution*, volume 13. Princeton University Press.
- Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, 16(2):97–159.